WORLD INTELLECTUAL PROPERTY ORGANIZATION





INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT) (51) International Patent Classification 7: (11) International Publication Number: WO 00/60940 A01N 25/04 A1 (43) International Publication Date: 19 October 2000 (19.10.00) (21) International Application Number: PCT/US00/09568 (81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CZ, DE, DK, DM, DZ, 10 April 2000 (10.04.00) EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, (22) International Filing Date: IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, (30) Priority Data: UG, UZ, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, 60/128,994 12 April 1999 (12.04.99) US BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, (71) Applicants: DOW AGROSCIENCES LLC [US/US]; 9330 NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA,

Zionsville Road, Indianapolis, IN 46268-1054 (US). THE DOW CHEMICAL COMPANY [US/US]; 2030 Dow Center, Midland, MI 48674 (US).

(72) Inventors: STROM, Robert, M.; 789 West Chippewa River Road, Midland, MI 48640 (US). PRICE, D., Claude; 2805 Canterbury Drive, Midland, MI 48667 (US). LUBETKIN, Steven, D.; 1135 Park Place, Zionsville, IN 46077 (US).

(74) Agent: WILLIS, Reid, S.; Intellectual Property, P.O. Box 1967, Midland, MI 48641-1967 (US).

Published

With international search report.

GN, GW, ML, MR, NE, SN, TD, TG).

Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: AQUEOUS DISPERSIONS OF AGRICULTURAL CHEMICALS

(57) Abstract

The biovailability of a pesticide can be increased by formulating the pesticide as a stable aqueous dispersion with a particle mean diameter of not greater than 500 nm. Such a formulation has the further advantage of substantially reducing or eliminating the need for organic solvents. Moreover, the stable aqueous dispersion provides a means of preparing a one part formulation of a plurality of pesticides which would be otherwise unstable in each other's presence.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FJ	Finland	LT	Lithuania	SK	Slovakia
ΑT	Austria	FR	France	LU	Luxembourg	SN	Senegal
ΑU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
ΑZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	1E	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Maurit ania	UG	Uganda
BY	Belarus	15	Iceland	MW	Malawi	US	United States of Americ
CA	Canada	lТ	Italy	МX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JР	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
СН	Switzerland	KG	Kyrgyzstan	NO	Norway	zw	Zimbabwe
Cl	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
· CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	K2	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	Ll	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

5

10

15

20

25

AQUEOUS DISPERSIONS OF AGRICULTURAL CHEMICALS

The present invention relates to an aqueous dispersion of particles having pesticidal activity. In particular, the present invention relates to dispersions of micron- or sub-micron-sized particles.

The efficient use of pesticides is often restricted somewhat by their inherent poor water-solubility. Generally, these water-insoluble pesticides can be applied to a site in three ways, 1) as a dust, 2) as a solution in an organic solvent or a combination of water and one or more organic solvents, or 3) as an emulsion that is prepared by dissolving the product in an organic solvent, then dispersing the solution in water. All of these approaches have drawbacks. Application of a dust poses a health hazard and is inefficient. Solutions and emulsions that require an organic solvent are undesirable since the solvent usually serves no other purpose but to act as a carrier for the product. As such, the solvent adds an unnecessary cost to the formulation. It would be an advantage in the art, therefore, to provide a pesticide formulation that eliminates the need for organic solvents, on the one hand, but exhibits optimal availability to the site to which it is applied.

The present invention provides a stable, aqueous, pesticidal dispersion comprising from 1 to 60 percent by weight of the aqueous dispersion of a pesticide having a water solubility of less than 0.1 percent, having a crystalline phase, and having a melting point sufficiently high so as not to melt during milling, and from 1 to 30 percent by weight based on the weight of the pesticide of a surface active agent, wherein the dispersion has a volume mean diameter particle size of not greater than 500 nanometers (nm).

In a second aspect, the present invention is a method of preparing a stable aqueous pesticidal dispersion comprising the steps of:

- a) mixing a pesticide with from 1 to 30 weight percent of a surface active agent and
 a sufficient amount of water to give an aqueous dispersion in the range of 1 to
 60 weight percent, based on the weight of water and the pesticide; and
- b) reducing the volume mean diameter particle size of the pesticide to not greater than 500 nm using mechanical means and a grinding medium.

The present invention addresses a need in the art by providing aqueous dispersions of submicron-sized particles having high bioavailability and no or minimal amounts of an ancillary organic solvent.

The stable aqueous dispersion of the pesticide can be prepared by wet milling an aqueous dispersion of the pesticide in the presence of grinding media and a surface active agent. As used herein, the term "a surface active agent" means one or more surface active agents. The preparation is carried out in such a manner so as to produce a dispersion of non-agglomerating or interacting particles having a volume median diameter of not greater than 500 nm, more preferably not greater than 400 nm, most preferably not greater than 200 nm, as measured by a Horiba model LA-700 particle size analyzer or an equivalent method. An example of such preparation can be found in U.S. Patent 5,145,684.

5

10

15

20

25

30

The pesticide is a solid at ambient temperature, and has a melting point that is sufficiently high that the pesticide does not melt through the milling process. The pesticide is preferably crystalline, and is substantially water insoluble. The term substantially water insoluble is used herein to refer to a pesticide that has a solubility in water of less than 0.1%, and most preferably less than 0.01%. It is understood that the water solubilities of many pesticides are pH-dependent, as a result of the functional groups they contain. Thus, pesticides with carboxylic acid groups or with sulfonamide or sulfonylurea groups, for example, may meet the solubility requirements at low pH but may be too soluble at high pH. The pH of the aqueous dispersion can be adjusted to ensure substantial insolubility of these pesticides.

Pesticides include herbicides, insecticides, and fungicides. Examples of classes of compounds that have herbicidal activity and meet the solubility, crystallinity and melting point requirements include, but are not restricted to, imidazolinones such as imazaquin, sulfonylureas such as chlorimuron-ethyl, triazolopyrimidine sulfonamides such as flumetsulam, aryloxyphenoxy propionates such as quizalofop ethyl, aryl ureas such as isoproturon and chlorotoluron, triazines such as atrazine and simazine, aryl carboxylic acids such as picloram, aryloxy alkanoic acids such as MCPA, chloroacetanilides such as metazachlor, dintroanilines such as oryzalin, pyrazoles such as pyrazolynate and diphenyl ethers such as bifenox.

Examples of classes of compounds that have insecticidal activity and meet the solubility, crystallinity and melting point requirements include, but are not restricted to, benzoyl ureas such as hexaflumuron, diacylhydrazines such as tebufenozide, carbamates such as carbofuran, pyrethroids such as *alpha*-cypermethrin, organophosphates such as phosmet, triazoles, and natural products such as spinosyns.

5

10

15

20

25

30

Examples of classes of compounds that have fungicidal activity and meet the solubility, crystallinity and melting point requirements include, but are not restricted to, morpholines such as dimethomorph, phenylamides such as benalaxyl, azoles such as hexaconazole, strobilurins such as azoxystrobin, phthalonitriles such as chlorothalonil and phenoxyquinolines such as quinoxyfen.

The surface active agent may be anionic, cationic, or nonionic, or combinations of cationic and nonionic or anionic and nonionic. Generally, and to a point, higher concentrations of surface active agent result in smaller particle size. Examples of suitable classes of surface active agents include, but are not limited to, anionics such as alkali metal fatty acid salts, including alkali metal oleates and stearates; alkali metal lauryl sulfates; alkali metal salts of diisooctyl sulfosuccinate; alkyl aryl sulfates or sulfonates, alkali metal alkylbenzene sulfonates, such as dodecylbenzene sulfonate; and alkali metal soaps; cationics such as long chain alkyl quaternary ammonium surfactants including cetyl trimethyl ammonium bromide; or nonionics such as ethoxylated derivatives of fatty alcohols, alkyl phenols, amines, fatty acids, fatty esters, mono-, di-, or triglycerides and various block copolymeric surfactants derived from ethylene oxide/propylene oxide. The concentration of the surface active agent is not less than 1%, and not more than 30% by weight, based on the total weight of the pesticide and the surface active agent. The surface active agent is preferably adsorbed onto the surface of the pesticide particle in accordance with U.S. Patent 5,145,684.

Examples of commercially available surface active agents include Atlox 4991 and 4913 surfactants (Uniqema), Morwet D425 surfactant (Witco), Pluronic P105 surfactant (BASF), Iconol TDA-6 surfactant (BASF), Kraftsperse 25M surfactant (Westvaco), Nipol 2782 surfactant (Stepan), Soprophor FL surfactant (Rhone-Poulenc), and Empicol LX 28 surfactant (Albright & Wilson).

The stable aqueous dispersions of the present invention can be prepared in the following manner. First, the pesticide is dispersed, preferably as a relatively coarse material having a particle size of less than 100 µm. Such a degree of coarseness can be achieved, for example, by sieving. The coarse material is dispersed in an aqueous solution containing the surface active agent and preferably an antifoaming agent such as Antifoam B (Dow Corning) to form a premix. The premix can then be ground mechanically to reduce further the particle size of the pesticide. The concentration of the pesticide is in the range of from 1% to 60%, more preferably to 30%, by weight based on the weight of the pesticide and the water. Attrition time can vary widely depending on the mechanical means used and the choice and concentration of surfactant.

10

15

20

25

30

Examples of mechanical means to reduce the size of the agricultural product include a ball mill, a roller miller, an attritor mill, a sand mill, a bead mill, and a Cowles type mixer. The grinding media used for particle size reduction preferably include substantially spheroidal shaped particles having an average size of less than 3 mm. Examples of suitable media material include ZrO stablized with magnesia, zirconium silicate, glass, stainless steel, polymeric beads, alumina, and titania, although nature of the material is not believed to be critical.

The particles are reduced in size at any suitable temperature where the agricultural product is stable. Typically, processing temperatures are not greater than the boiling point of water and below the melting point of the solid, but ambient temperature is preferred. The final volume mean diameter particle size is less than 500 microns, more preferably less than 400 nm, and most preferably less than 200 nm.

The stable aqueous dispersions are preferably diluted with water, then applied to a site in a pesticidally effective amount.

One of the advantages of the stable aqueous dispersion of the present invention is that it provides a means to prepare one-part formulations of different pesticides which are not only compatible with each other, but incompatible or unstable in each other's presence as well. For example, it may be desirable to combine a certain pesticide with a certain herbicide for a particular application but for the fact that the two (in solution, for example) react with each other faster than they can be applied to the desired site. However, in a stable aqueous dispersion, these different and incompatible pesticides can coexist, at least

temporarily, since they are shielded from each other from reacting rapidly, so that an end user can mix the incompatible pesticides together and apply them to a site before their efficacy is significantly diminished.

The fine particle size suspensions prepared according to the methods of the present invention, have enhanced biological efficacy as a result of their increased surface area as compared to suspensions of greater particle size. Enhanced biological efficacy is particularly important in cases where pesticide solubility is very low and thus, biological availability is restricted.

The following examples are for illustrative purposes only and are not intended to limit
the scope of the invention. All percentages are by weight unless otherwise specified. All
particle sizes (measured using a Horiba model LA-700 particle size analyzer) are in units of
volume mean diameter.

Example 1 - Preparation of a Stable Aqueous Nano-Dispersion of a Triazole

5

15

20

1-Methyl- 3-(2-fluoro-6-chlorophenyl)-5-(3-methyl-4-bromothien-2-yl)-1H-1,2,4-triazole (1 g) was placed in a 20 mL polyethylene scintillation vial along with an aqueous phase (9 g) that was prepared by combining Pluronic P105 surfactant (3.33%), Morwet D425 surfactant (2.22%), deionized water (93.33%) and Antifoam B antifoaming agent (1.11%). About 10 g of 1/8" diameter stainless steel ball bearings were added, and the vial was then sealed, and placed a Retsch Model MM 2000 mill. The sample was milled for 90 minutes and the particle size (pre-mill) was measured to be 1520 nm. A portion of the pre-mill (1 g) was removed and added to a fresh vial, at which time 1 drop of the antifoaming agent was added. Milling was continued for 3 hours, after which the particle size was determined to be 183 nm.

<u>Example 2</u> – Preparation of a Stable Aqueous Nano-Dispersion Using Polymeric Beads

Example 1 was repeated except that styrene/divinylbenzene copolymer beads (475 μm diameter) were used instead of stainless steel beads. After the pre-mill was subjected to 120 minutes of additional milling, the particle size was determined to be 148 nm.

<u>Example 3</u> – Preparation of a Stable Aqueous Nano-Dispersion Using Alternative Surfactant Package

Example 1 was repeated except that the aqueous phase was prepared using Atlox 4991 surfactant (6%), Atlox 4913 surfactant (12%), deionized water (78%) and Antifoam B (4%). After the pre-mill was subjected to 270 minutes of additional milling, the particle size was determined to be 163 nm.

5 Example 4 - Preparation of a Stable Aqueous Nano-Dispersion of Spinosad

In this example, Spinosad insecticide was milled. Again, Example 1 conditions were used except that the dispersion was prepared using the surfactant package described in Example 3. After the pre-mill was subjected to 195 minutes of additional milling, the particle size was determined to be 297 nm.

10 Example 5 - Preparation of a Stable Aqueous Nano-Dispersion of Spinosad Using a Different Surfactant Package

The conditions of Example 1 were repeated except the Spinosad was used instead of the triazole. After the pre-mill was subjected to 315 minutes of additional milling, the particle size was found to be 281 nm.

15 Example 6 - Preparation of a Stable Aqueous Nano-Dispersion of Atrazine Herbicide

20

Atrazine herbicide (1 g) was placed in a 20 mL polyethylene scintillation vial along with an aqueous phase (9 g) prepared by combining Iconol TD-6 surfactant (0.5 g) deionized water (8 g) and Antifoam B (0.5 g). About 10 g of 1/8" stainless steel ball bearings were added, and the vial was sealed and placed on the mill. The particles were milled for 30 minutes, after which time a portion of the pre-mill (1 g) was removed and placed in a fresh vial along with fresh ball bearings (10 g) and 1 drop of Antifoam B. The vial was sealed and subjected to milling for an additional 105 minutes. Then Nipol 2782 surfactant (0.2 g) was added and milling was continued for an additional 2 hours. The particle size was measured to be 167 nm.

25 <u>Example 7</u> - Preparation of a Stable Aqueous Nano-Dispersion of Epoxiconazole Herbicide

Epoxiconazole was placed in a 20 mL polyethylene scintillation vial along with an aqueous phase prepared from Soprophor FL surfactant (0.2 g), Empicol LX 28 (0.3 g), deionized water (8.45 g), and Antifoam B agent (0.05 g). About 10 g of 1/8" steel ball

10

bearings were added, and the vial was sealed and placed on the mill for 30 minutes, after which time, a portion of the pre-mill (1 g) was removed and placed in a fresh vial along with fresh ball bearings (10 g) and 1 drop of Antifoam B. The vial was sealed and subjected to milling for an additional 405 minutes. The particle size was measured to be 314 nm.

A correlation between particle and biological efficacy as demonstrated LC 50 is shown in the Table I below.

Table I

Two spotted :	spider mite
VMD/ nm	LC
0.404	15
0.372	11
0.332	7.6
0.163	4

As shown by the table, the amount required to kill 50% of the spider mites is substantially reduced at smaller VMD.

WHAT IS CLAIMED IS:

5

10

15

20

1. A stable, aqueous pesticidal dispersion comprising from 1 to 60 percent by weight of the aqueous dispersion of a pesticide having a water solubility of less than 0.1 percent, having a crystalline phase, and having a melting point sufficiently high so as not to melt during milling, and from 1 to 30 percent by weight based on the weight of the pesticide of a surface active agent, wherein the dispersion has a volume mean diameter of not greater than 500 nm.

- 2. The pesticidal dispersion of Claim 1 wherein the pesticide is a herbicide selected from the group consisting of imidazolinones, sulfonylureas, triazolopyrimidine sulfonamides, aryloxyphenoxy propionates, aryl ureas, triazines, aryl carboxylic acids, aryloxy alkanoic acids, chloroacetanilides, dintroanilines, pyrazoles, and diphenyl ethers.
- 3. The pesiticidal dispersion of Claim 1 wherein the pesticide is an insecticide selected from the groups consisiting of benzoyl ureas, diacylhydrazines, carbamates, pyrethroids, organophosphates, triazoles, and natural products.
- 4. The pesticidal dispersion of Claim 1 wherein the pesticide is a fungicide selected from the group consisting of morpholines, phenylamides, azoles, strobilurins, phthalonitriles, and phenoxyquinolines.
- 5. The pesticidal dispersion of any of Claims 1 to 4 wherein the surface active agent is selected from the group consisting of alkali metal fatty acid salts, polyoxyethylene nonionics, alkali metal lauryl sulfates, quaternary ammonium surfactants, alkali metal alkylbenzene sulfonates, and alkali metal soaps.
- 6. The pesticidal dispersion of any of Claims 1 to 5 which has a volume mean diameter of not greater than 200 nm.
- 7. A method for controlling pests comprising the step of diluting the stable aqueous dispersion of any of Claims 1 to 6 with water and applying to a site a pesticidally effective amount of the diluted dispersion.

5

10

8. A method of preparing a stable aqueous pesticidal dispersion comprising the steps of:

- a) mixing a pesticide with from 1 to 30 weight percent of a surface active agent and a sufficient amount of water to give an aqueous dispersion in the range of 1 to 60 weight percent, based on the weight of water and the pesticide; and
- b) reducing the volume mean diameter particle size of the pesticide to not greater than 500 nm using mechanical means and a grinding medium.
- 9. The method of Claim 8 wherein the mechanical means is a ball mill, a roller mill, an attritor mill, a sand mill, a bead mill, or a Cowles type mixer.
- 10. The method of either of Claims 8 or 9 wherein the grinding media are particles selected from the group consisting of ZrO stabilized with magnesia, zirconium silicate, glass, stainless steel, polymeric beads, alumina, and titania.

PCT/US 00/09568

A. CLASSIFICATION OF SUBJECT MATTER IPC 7 A01N25/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols) IPC 7 A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS, CHEM ABS Data

C. DOCUMENTS	CONSIDERED .	TO BE RELEVANT

WO 99 48359 A (MULQUEEN PATRICK JOSEPH; BANKS GRAHAM (GB); DIXON KEITH RODERICK () 30 September 1999 (1999-09-30) page 1 -page 2 page 5 -page 10 claims 2-7,20	1-10
WO 95 07614 A (DOWELANCO; MULQUEEN PATRICK JOSEPH (GB); BANKS GRAHAM (GB); LUBETK) 23 March 1995 (1995-03-23) page 1, paragraph 1 page 20, paragraph 3 examples 21-23	1-10
EP 0 499 299 A (STERLING WINTHROP INC) 19 August 1992 (1992-08-19) cited in the application page 2, line 50 -page 3, line 13; examples	1-10
	page 20, paragraph 3 examples 21-23 EP 0 499 299 A (STERLING WINTHROP INC) 19 August 1992 (1992-08-19) cited in the application

	-/
Further documents are listed in the continuation of box C.	Patent family members are listed in annex.
Special categories of cited documents: 'A' document defining the general state of the art which is not considered to be of particular relevance 'E' earlier document but published on or after the international filling date 'L' document which may throw doubts on priority claim(s) or which is cited to establish the publication cate of another citation or other special reason (as specified) 'O' document referring to an oral disclosure, use, exhibition or other means 'P' document published prior to the international filing date but later than the priority date claimed	 'T' later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention 'X' document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone 'Y' document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. '8" document member of the same patent family
Date of the actual completion of the international search 2 August 2000	Date of mailing of the international search report 11/08/2000
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016	Authorized officer Bertrand, F

1

Inter onal Application No PCT/US 00/09568

•		1 101/03	00/09568
C.(Continu	etion) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category °	Citation of document, with indication, where appropriate, of the relevant passages	;:	Relevant to claim No.
Y	GB 2 267 825 A (DOWELANCO) 22 December 1993 (1993-12-22) the whole document		1-10
X	US 4 832 950 A (TAKAYA SUSUMU ET AL) 23 May 1989 (1989-05-23) column 1 -column 3; examples		1-10
(EP 0 683 979 A (AMERICAN CYANAMID CO) 29 November 1995 (1995-11-29) the whole document		1-10
1	US 4 430 336 A (RENTZEA COSTIN ET AL) 7 February 1984 (1984-02-07) column 17, line 62 -column 18, line 4		1-10
		.•	

1

Inte ional Application No PCT/US 00/09568

		7			L	L1/US	00/09568
	atent document d in search report		Publication date	P 	atent family member(s)		Publication date
WO	994835 9	A	30-0 9- 199 9	AU	2946299	A	18-10-1999
WO	9507614	Α	23-03-1995	AU	691835	В	28-05-1998
				ΑU	7835594		03-04-1995
				BR	9407501	Α	25-06-1996
				CA	2171848	Α	23-03-1995
				ΕP	0719086		03-07-1996
				HU	74022	A,B	28-10-1996
				ΙL	110993		15-07-1998
				US	607498 6		13-06-2000
				ZA 	94 07147 	A	15-03-1996
EP	04992 99	Α	19-08-1992	US	5145684		08-09-1992
				AT	184202		15-09-1999
				AU AU	642066 1014592		07-10-1993 30-07-1992
				AU	654836		24-11-1994
				AU	1014792		30-07-1992
				CA	2059431		26-07-1992
				CA	2059432		26-07-1992
				DE	69229925		14-10-1999
				DE	69229925	T	17-02-200 0
				EP	0498482		12-08-1992
				ES	2139586		16-02-2000
				FI	920321		26-07-1992
				FI	920322		26-07-1992
			•	HU	62462		28-05-1993
				HU	60635		28-10-1992
				IL IL	10075 4 100755		16-10-1996
	•			JP	4317053		08-12-1995 09-11-1992
				JP	4295420		20-10-1992
				KR	200061		15-06-199 9
				MX	9200291		01-10-1992
				MX	9200292		01-10-1992
				NO	920333		27-07-1992
				NO	303668	В	17-08-1998
				NZ	241361		25-06-1993
				NZ	241362		25-06-1993
				SG	55104		21-12-1998
				RU	2074002		27-02-1997
				RU	2066553		20-09-1996
				U\$	5451393		19-09-1995
			i	US US	549468 3 555216 0		27-02-1996
				US	5399363		03-09-1996 21-03-1995
				US	531876 7		07-06-1994
GB	22 6 78 25	Α	22-12-1993	NONE			
US	4832950	Α	23-05-1989	JP	149979 0		29-05-198 9
				JP	61057503		24-03-1986
				JP	63052002		17-10-1988
				JP	1011610	В	27-02 - 198 9
				JP	15368 17	C	21-12-1989
				JP	61151111	Α	09-07-1986
						A C	

Inter: Inal Application No PCT/US 00/09568

Patent document	Publication		Patent family	Publication
cited in search report	date		member(s)	Publication date
US 4832950 A		JP	61152627 A	11-07-1986
		AT	6915 8 T	15-11-1991
		DE	358459 5 A	12-12-1991
		EP	0173259 A	05-03-1986
		HK	415 93 A	07-05-1993
		PH	230 96 A	10-04-198 9
		S G	14093 G	16-04-1993
EP 0683979 A	29-11-1995	US	549684 5 A	05-03-1996
		AT	18204 8 T	15-07-199 9
<u>.</u> •		AU	70794 4 B	22-07-1999
		AÜ	20196 95 A	30-11-1995
		BG	625 06 B	31-01-2000
		BG	996 65 A	29-02-1996
		BR	9502522 A	
				09-04-1996
		CA	2149969 A	26-11-1995
		CN	1114134 A	03-01-1996
		CZ	9501259 A	13-12-1995
		DE	6951072 9 D	19-08-1999
		DE	6951072 9 T	04-11-199 9
		EG	203 92 A	28-02-199 9
		ES	2133672 T	16-09-1999
		GR	30 31104 T	31-12-1999
	i .	HU	73158 A	28-06-1996
		JP	8040809 A	13-02-1996
		NO	952061 A	27-11-1995
		NZ	272184 A	24-03-1997
		PL	3087 37 A	27-11-1995
		RO	112684 A	30-12-1997
		SI	683979 T	31-10-1999
•		SK	68795 A	
		TR		06-12-1995
			285 83 A	04-11-1996
		ZA 	95042 15 A	22-01-1996
US 4430336 A	07-02-1984	DE	2948704 A	11-06-1981
		AT	2510 T	15-03-1983
		AU	650318 0 A	11-06-1981
		BR	80076 90 A	09-06-1981
		CA	1153002 A	30-08-1983
		CS	215143 B	30-07-1982
		DE	3062078 D	24-03-1983
		DK	515180 A	05-06-1981
		EP	0029996 A	10-06-1981
		HU	185622 B	28-03-1985
		JE	50726 B	25-05-1985 25-06-1986
		IL	61515 A	29-02-1984
		JP	56092851 A	27-07-1981
		NZ	195740 A	07-12-1982
		PL	228213 A	07-08-1981
		C 1 1	1000005 4	07 10 1000
		SU	10600 95 A	07-12-1983